

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 31/66	A1	(11) International Publication Number: WO 96/39151 (43) International Publication Date: 12 December 1996 (12.12.96)
(21) International Application Number: PCT/US96/08398 (22) International Filing Date: 3 June 1996 (03.06.96) (30) Priority Data: 08/471,466 6 June 1995 (06.06.95) US (60) Parent Application or Grant (63) Related by Continuation US 08/471,466 (CON) Filed on 6 June 1995 (06.06.95) (71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): DAIFOTIS, Anastasia, G. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). YATES, Ashley, J. [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). (74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		(81) Designated States: AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: BISPHOSPHONATES PREVENT BONE LOSS ASSOCIATED WITH IMMUNOSUPPRESSIVE THERAPY (57) Abstract <p>Bisphosphonate, particularly alendronate, can prevent or treat bone loss associated with immunosuppressive therapy, whether or not the immunosuppressive therapy is associated with an organ transplant.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

TITLE OF THE INVENTION

BISPHOSPHONATES PREVENT BONE LOSS ASSOCIATED WITH
IMMUNOSUPPRESSIVE THERAPY

5 SUMMARY OF THE INVENTION

This invention relates to the use of bisphosphonates,
particularly alendronate, to prevent bone loss associated with
immunosuppressive therapy, and in particular when such therapy is used
in conjunction with organ transplantation.

10

BACKGROUND OF THE INVENTION

Patients suffering from various medical conditions which
require an organ or bone marrow transplant, need a variety of drugs in
order to suppress the body's tendency to reject the organ. This
15 generally requires that the patient take one or more immunosuppressive
agents, such as cyclosporine or the like, often in combination with
adrenal corticosteroids, such as methylprednisolone. Unfortunately, the
combination of the underlying condition, immobility or decreased
mobility, and drug therapy causes these patients to experience a high
20 degree of bone loss.

Further, various immunosuppressive agents are being tried
as therapeutic agents in treating various conditions which do not
necessarily involve organ transplantation, such as in rheumatoid
arthritis, psoriasis, inflammatory bowel disease and nephrotic
25 syndrome. These patients also are at high risk for bone loss.

It would be desirable to be able to combat or prevent bone
loss in patients who are undergoing organ transplants or receiving
immunosuppressive therapy in association with an organ transplant or
other underlying medical condition.

30

DETAILED DESCRIPTION OF THE INVENTION

It has been found in accordance with this invention that
bisphosphonates can prevent and treat bone loss associated with
immunosuppressive therapy when administered either in either a

- 2 -

prophylactically or therapeutically effective amount. In particular, alendronate (4-amino-1-hydroxybutylidene-1,1-bisphosphonate) or a pharmaceutically effective salt thereof, can prevent and treat bone loss associated with organ transplants when administered either in either a
5 prophylactically or therapeutically effective amount.

A further aspect of this invention is to prevent or treat bone loss associated with immunosuppressive therapy, regardless of whether the therapy accompanies organ transplantation by administration of an effective amount of a bisphosphonate selected from the group consisting
10 of: alendronate, etidronate (1-hydroxy-ethidene-bisphosphonic acid), pamidronate (3-amino-1-hydroxypropylidene-1,1-diphosphonate), risedronate (2-(3-pyridinyl)-1-hydroxyethylidene-bisphosphonic acid), clodronate (dichloromethylene-bisphosphonic acid), tiludronate (chloro-4-phenylthiomethylidene-bisphosphonic acid), ibandronic acid (1-
15 hydroxy-3(methylpentylamino)-propylidene-bisphosphonic acid, and pharmaceutically acceptable salts of any of the foregoing, and mixtures of any of the acids and any of the salts. All of the foregoing compounds are well known in the art.

Generally the patient undergoing immunosuppressive
20 therapy in accordance with this invention will be receiving cyclosporine or a similar drug. In addition, the patient may also be receiving prednisone or other corticosteroid.

As used throughout the specification and claims, the
25 following definitions apply:

"Prophylactically effective amount"--the amount of alendronate needed to prevent or lessen the severity of bone loss related to immunosuppressive therapy, regardless of whether the immunosuppressive therapy is accompanied by organ transplantation.

30 "Therapeutically effective amount"--the amount of alendronate needed to treat bone loss related to immunosuppressive therapy, regardless of whether the immunosuppressive therapy is accompanied by organ transplantation.

- 3 -

In a preferred aspect of this invention, the patient will receive alendronate. Alendronate may be prepared according to any of the processes described in U.S. Patents 5,019,651, 4,992,007, and U.S. Application Serial No. 08/286,151, filed August 4, 1994, each of which is hereby incorporated by reference. The pharmaceutically acceptable salts of alendronate include salts of alkali metals (e.g., Na, K), alkaline earth metals (e.g. Ca), salts of inorganic acids, such as HCl and salts of organic acids such as citric acid and amino acids. Sodium salt forms are preferred, particularly the monosodium salt trihydrate form.

Many of the bisphosphonate compounds of the present invention can be administered in oral dosage forms such as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, paste, tinctures, suspensions, syrups, and emulsions. Likewise they may be administered in an intravenous (bolus or infusion), intraperitoneal, subcutaneous, or intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts. An effective but non-toxic amount of the bisphosphonate compound desired can be used to prevent bone loss.

The dosage regime utilizing the claimed method is selected in accordance with a variety of factors including type, age, weight, sex, and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt thereof employed. An ordinarily skilled physician or clinician can readily determine and prescribe the effective amount of the drug required to prevent and or treat bone loss.

Oral dosages of the present invention when alendronate is the bisphosphonate will range from between 0.05 mg per kg of body weight per day (mg/kg/day) to about 1.0 mg/kg/day. Preferred oral dosages in humans may range from daily total dosages of about 2.5-50 mg/day over the effective treatment period, and a preferred amount is 5, 10 or 20 mg/day.

Alendronate may be administered in a single daily dose or in a divided dose. It is desirable for the dosage to be given in the

- 4 -

absence of food, preferably from about 30 minutes to 2 hours prior to a meal, such as breakfast to permit adequate absorption.

In the methods of the present invention, the active ingredient is typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier materials") suitably selected with respect to the intended form of administration, i.e. oral tablets, capsules, elixirs, syrups and the like and consistent with conventional pharmaceutical practices. For example, for oral administration in the form of a tablet or capsule, the active ingredient can be combined with an oral, non-toxic, pharmaceutically acceptable inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, mannitol, sorbitol, croscarmellose sodium and the like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture of active ingredient(s) and inert carrier materials. Suitable binders may include starch, gelatin, natural sugars such as glucose, anhydrous lactose, free-flow lactose, beta-lactose, and corn sweeteners, natural and synthetic gums, such as acacia, tragacanth or sodium alginate, carboxymethyl cellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. A particularly preferred tablet formulation for alendronate is that described in U.S. Patent 5,358,941, which is hereby incorporated by reference.

The compounds used in the instant method may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran co-polymer, polyhydroxypropyl-methacrylamide and the like.

Patients undergoing immunosuppressive therapy may be male or female of any age. Women may be pre- or post-menopausal.

- 5 -

The following non-limiting examples are presented to better illustrate the invention.

EXAMPLE 1

5

Alendronate for the prevention and treatment of cyclosporine-induced bone loss

220 men and women are enrolled in a clinical study to assess the effectiveness of alendronate to prevent and treat bone-loss associated with immunosuppressive therapy. All subjects are receiving an organ transplant; the majority are receiving a heart, lung or liver transplant. The patients are randomized into 5 groups which receive either placebo, 2.5, 5, 10, or 20 mg/day alendronate for one year, beginning within one week post transplant. In addition to standard amounts of cyclosporine and steroid such as prednisone, all patients also receive 1000 mg per day calcium and 250 IU per day Vitamin D.

15

Spine and hip bone mineral densities are monitored, and all incidences of fractures are recorded.

After one year, patient receiving alendronate (at any dose) have a statistically significantly higher spine and hip BMD than placebo patients, and have experienced less fractures. This result is observed in both strata: Those with low starting BMD are seen to gain BMD. Those whose starting BMD is not low are observed to retain BMD. Thus alendronate prevents and treats bone loss associated with immunosuppressive therapy.

20
25

- 6 -

WHAT IS CLAIMED IS:

1. A method of treating or preventing bone associated with immunosuppressive therapy comprising: administering an effective amount of a bisphosphonate selected from the group consisting of:
alendronate, etidronate (1-hydroxy-ethidene-bisphosphonic acid), pamidronate (3-amino-1-hydroxypropyldiene-1,1-diphosphanate), risedronate (2-(3-pyridinyl)-1-hydroxyethylidene-bisphosphonic acid), clodronate (dichloromethylene-bisphosphonic acid), tiludronate (chloro-4-phenylthiomethylidene-bisphosphonic acid), ibandronic acid (1-hydroxy-3(methylpentylamino)-propylidene-bisphosphonic acid, pharmaceutically acceptable salts of any of the foregoing, and mixtures thereof to a patient undergoing immunosuppressive therapy.
2. A method according to Claim 1 wherein the patient is receiving immunosuppressive therapy in association with an organ transplant.
3. A method according to Claim 2 comprising administering alendronate or a pharmaceutically acceptable salt thereof.
4. A method according to Claim 3 wherein the alendronate is in the form of monosodium salt trihydrate.
5. A method according to Claim 4 wherein the alendronate is administered in a prophylactically effective amount.
6. A method according to Claim 5 wherein the alendronate is administered orally.
7. A method according to Claim 5 wherein the alendronate is administered in a dose of from 2.5 to 50 mg per day.

- 7 -

8. A method according to Claim 7 wherein the alendronate is administered in a dose of 5 mg, 10 mg, or 20 mg per day.

5 9. A method of treating bone loss associated with immunosuppressive therapy comprising administering a therapeutically effective amount of alendronate or a pharmaceutically acceptable salt thereof to a patient undergoing immunosuppressive therapy.

10 10. A method according to Claim 9 wherein the alendronate is in the form of monosodium salt trihydrate.

11. A method according to Claim 10 wherein the alendronate is administered orally.

15 12. A method according to Claim 10 wherein the alendronate is administered in a dose of from 2.5 to 50 mg per day.

20 13. A method according to Claim 12 wherein the alendronate is administered in a dose of 5 mg, 10 mg, or 20 mg per day.

14. A method according to Claim 9 wherein the immunosuppressive therapy is associated with an organ transplant.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/08398

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/66

US CL :514/102, 108

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/102, 108

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS:

TERMS SEARCHED: BISPHOSPHONATES AND IMMUNOSUPPRESSIVE AND BONE AND ALENDRONATE.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, E	US, A, 5,536,716 (CHEN et al.) 16 July 1996, column 36, lines 49-64.	1-14



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*g* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

22 AUGUST 1996

Date of mailing of the international search report

17 SEP 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

THEODORE J. CRIARES

Telephone No. (703) 308-1235